



P: 1300 688 522  
E: info@nutripath.com.au  
A: PO Box 442 Ashburton VIC 3142

**TEST PATIENT**

GUa d'Y'HYghBUa Y  
Sex : :  
DUH Y Collected : 00-00-0000  
111 H9GH ROAD TEST SUBURB  
**@AB =8: 00000000** UR#:0000000

**TEST PHYSICIAN**

DR JOHN DOE  
111 CLINIC STF 99H  
7@=B=7'GI 6I F 6'J =7'' \$\$\$

**GIT ASSESSMENTS**

STOOL, SPOT

**Pancreatic Elastase 1**

Result	Range	Units
>500	> 200	ug/g

**Digestive Markers Comment**

**PANCREATIC ELASTASE:** Normal exocrine pancreatic function.

Pancreatic Elastase reflects trypsin, chymotrypsin, amylase and lipase activity. This test is not affected by supplements of pancreatic enzymes.

Healthy individuals produce on average 500 ug/g of PE-1. Thus, levels below 500 ug/g and above 200 ug/g suggest a deviation from optimal pancreatic function.

The clinician should therefore consider digestive enzyme supplementation if one or more of the following conditions is present:

Loose watery stools, Undigested food in the stools, Post-prandial abdominal pain, Nausea or colicky abdominal pain, Gastroesophageal reflux symptoms, Bloating or food intolerance.

Tests ordered: PE-1,IMPEI

(\*) Result outside normal reference range

(H) Result is above upper limit of reference rang