



TEST PATIENT

GUa d'Y'HYgh'BUa Y
 Sex : :
 DUHY Collected : 00-00-0000
 111 H9GH'ROAD TEST SUBURB
@AB =8: 0000000 UR#:0000000

TEST PHYSICIAN

DR JOHN DOE
 111 CLINIC STF 99H
 7@-B =7 'GI 6I F 6'J =7'' \$\$\$

P: 1300 688 522
 E: info@nutripath.com.au
 A: PO Box 442 Ashburton VIC 3142

ENDOCRINOLOGY SALIVA

SALIVA	Result	Range	Units	
Male Hormone Profile-Extensive				
DHEAS.	18.9	5.0 - 30.0	nmol/L	
Androstenedione...	1.4	1.0 - 10.7	nmol/L	
Testosterone.	389.0	100.0 - 720.0	pmol/L	
Salivary DHT	2.8	2.0 - 10.0	pg/mL	
Estradiol (E2)	10.0		pmol/L	
Estrone (E1)	18.0	9.6 - 20.0	pg/mL	
Testosterone/DHT Ratio	138.9	10.0 - 350.0	RATIO	
Androstenedione/E1 Ratio	0.08	0.05 - 1.10	RATIO	



TEST PATIENT

GUa d'Y'HYghBUa Y
 Sex : :
 DUHY Collected : 00-00-0000
 111 H9GH ROAD TEST SUBURB
@AB =8: 0000000 UR#:0000000

TEST PHYSICIAN

DR JOHN DOE
 111 CLINIC STF 99H
 7@B=7'GI 6I F6'J =7" \$\$\$

P: 1300 688 522
 E: info@nutripath.com.au
 A: PO Box 442 Ashburton VIC 3142

Saliva Hormone Comments

SALIVARY HORMONE REFERENCE RANGES: (NOT ON HRT - BASELINE)

	E2	E1	E3	Progesterone	DHEAS
FEMALE					
Follicular	<18	9.6-20	15-29	<318	
Mid-Cycle	11-29	9.6-20	15-29	-	
Luteal	<18	9.6-20	15-29	318-1590	
Post Men.	<6	9.6-20	1-41	<159	<6.5
Premenopausal, no oral contraceptives					2.5-25.0
Premenopausal, with oral contraceptives					2.0-8.0
MALE					
	<6	9.6-20	16-25	<159	5.0-30.0

TARGET REFERENCE RANGES: (ON HRT - 24hr post last dose)

	E2	E1	E3	Progesterone	Testosterone Age Dpndt
Oral	7-73	-	69-139	318-1590	
Patch	4-18	-	-	-	
Cream/Gel	37-184	-	1040-1734	3180-31797	F: 277-867 M: 347-1734

SALIVA ESTRONE (E1) is produced primarily from androstenedione originating from the gonads or the adrenal cortex. In premenopausal women, more than 50% of the E1 is secreted by the ovaries. In prepubertal children, men and non-supplemented postmenopausal women, the major portion of E1 is derived from peripheral tissue conversion of androstenedione. Interconversion of E1 and E2 also occurs in peripheral tissue. Bioassay data indicate that the estrogenic action is much less than E2. E1 is a primary estrogenic component of several pharmaceutical preparations, including those containing conjugated and esterified estrogens. In premenopausal women E1 levels generally parallel those of E2. After menopause E1 levels increase, possibly due to increased conversion of androstenedione to E1.

SALIVA E1 level is within range for a male, indicating minimal aromatisation of androgens to E1.

ELEVATED E2 LEVEL:

Saliva E2 level is elevated for a male and suggestive of aromatisation of androgens to estrogens. Suggest using 5% transdermal Chrysin and / or 50mg Zinc.

SALIVA DHEAs level is adequate and within range.

SALIVA FREE TESTOSTERONE level is adequate and within range.