



P: 1300 688 522
E: info@nutripath.com.au
A: PO Box 442 Ashburton VIC 3142

TEST PATIENT

GUa d`Y`HYghBUa Y
Sex : :
DUHY Collected : 00-00-0000
111 H9GH`ROAD TEST SUBURB
@AB =8: 0000000 UR#:0000000

TEST PHYSICIAN

DR JOHN DOE
111 CLINIC STF 99H
7@B=7`GI 6I F6`J =7` \$\$\$

MICROBIOLOGY

STOOL, SPOT Result Range Units

Faecal Virus Profile

ENTERIC VIRAL PATHOGENS PROFILE:

Rotavirus: DETECTED
Adenovirus: Not Detected
Astrovirus: Not Detected
Norovirus: DETECTED

Comments:

Several viruses are known to cause gastroenteritis. On a global basis viral enteric disease accounts for 60% of illness in developing countries and 40% in developed countries. Enteric viral infections are seen more frequently in infants and children and somewhat lesser in adults.

Rotavirus DETECTED:

Rotavirus is the leading cause of severe diarrhoea in infants and young children.

Norovirus DETECTED:

Noroviruses are the leading cause of gastroenteritis, including foodborne outbreaks. It can be transmitted by consuming contamination food or drinks, touching contaminated objects then putting hand/fingers in mouth or direct contact with infected individual e.g. sharing food, drink, eating utensils.

